

Person-centered Critical Care Nursing (PCCN)

- English ver. -



Dong-A university, College of Nursing,
The Person Centered Critical Care Team

601, College of Nursing, Dong-A University, 32, Daesingongwon-ro, Seo-gu, Busan, Republic of Korea

Citation

- ✓ Kang J, Cho YS, Jeong YJ, Kim SG, Yun S, & Shim M. (2018). Development and Validation of a Measurement to Assess Person-centered Critical Care Nursing. *Journal of Korean Academy of Nursing*, 48(3), 323-334.

Instructions

- ✓ 15 items, 4 factors (Item No. 1~4=Compassion, 5~8=Individuality, 9~12=Respect, 13~15=Comfort).
- ✓ A 5-point Likert-type scale was used to measure PCCN (1 = Not at all ~ 5= Extremely).
- ✓ A high score means a high degree of person-centered critical care nursing performance by intensive care unit nurses.

Copyright© 2021 by The Person Centered Critical Care Team.
This measurement tool cannot be copied without permission.

Measurement to Assess Person-centered Critical Care Nursing (English version)

The following questions are about person-centered Critical Care Nursing by nurses in the intensive care unit. Please fill in the V mark in the relevant column of each question.

1: Not at all 2: A little bit 3: Moderately 4: Quite a bit 5: Extremely

No.	Items	1	2	3	4	5
1	Condole patient and family with words and actions.	1	2	3	4	5
2	Try to empathize with the situation of the patient and family.	1	2	3	4	5
3	Try to make therapeutic contact with the patient and family (e.g., holding hands).	1	2	3	4	5
4	Talk to the patient and family about everyday topics (e.g., news, hobbies, interests).	1	2	3	4	5
5	Allow the patient to bring in the desired item (e.g., doll, religious article, photo, pillow).	1	2	3	4	5
6	Allow the patient to enjoy desired entertainment activities (e.g., listening to music, watching videos, reading).	1	2	3	4	5
7	In addition to the scheduled time, allow family visits if necessary.	1	2	3	4	5
8	Allow the family to participate in patient care (e.g., personal hygiene, feeding).	1	2	3	4	5
9	Talk to the unconscious patient as if he/she can listen to you.	1	2	3	4	5
10	Try non-verbal communication with the patient who can not speak.	1	2	3	4	5
11	Allow the patient to urinate or defecate in the way he/she wants, to the extent possible.	1	2	3	4	5
12	Consider other patients in the vicinity when CPR or death is ongoing.	1	2	3	4	5
13	Avoid unnecessary noise (e.g., chattering of medical staff, alarms).	1	2	3	4	5
14	Try to reduce unpleasant odor in the intensive care unit.	1	2	3	4	5
15	Adjust lighting according to patients' preference at night.	1	2	3	4	5

This measurement tool cannot be copied without permission.